

# Improved $q$ -exponential and $q$ -trigonometric functions

Jan L. Cieśliński\*

Uniwersytet w Białymstoku, Wydział Fizyki  
ul. Lipowa 41, 15-424 Białystok, Poland

## Abstract

We propose a new definition of the  $q$ -exponential function. Our  $q$ -exponential function maps the imaginary axis into the unit circle and the resulting  $q$ -trigonometric functions are bounded and satisfy the Pythagorean identity.

*MSC 2010:* 05A30; 33B10; 39A13

*Key words and phrases:* quantum calculus,  $q$ -calculus,  $q$ -exponential function,  $q$ -trigonometric functions, Cayley transform,

---

\*e-mail: [janek@alpha.uwb.edu.pl](mailto:janek@alpha.uwb.edu.pl)

# 1 Introduction

The quantum calculus ( $q$ -calculus) is an old, classical branch of mathematics, which can be traced back to Euler and Gauss [11, 21] with important contributions of Jackson a century ago [18, 19]. In recent years there are many new developments and applications of the  $q$ -calculus in mathematical physics, especially concerning special functions [1, 8, 12, 14] and quantum mechanics [4, 5, 25, 13, 10, 23, 29]. Many papers were devoted to various approaches to  $q$ -deformations of elementary functions, including exponential and trigonometric functions [2, 3, 7, 15, 22, 24, 26, 27, 28].

In this paper we propose new definitions of the  $q$ -exponential function and  $q$ -trigonometric functions. These results are motivated by recent developments in the time scales calculus, where new exponential, hyperbolic and trigonometric function have been defined [9]. The concept of time scales unifies difference and differential calculus [16]. The  $q$ -calculus can be considered as a calculus on a special time scale (see, e.g., [6]).

The functions presented in this paper have better qualitative properties than standard  $q$ -exponential and  $q$ -trigonometric functions. In order to discuss and compare these properties we begin with a short summary of the classical results, usually following the textbook [20].

In the standard approach to the  $q$ -calculus two exponential function are used:

$$e_q^z = \sum_{n=0}^{\infty} \frac{z^n}{[n]!} , \quad E_q^z = \sum_{n=1}^{\infty} \frac{z^n}{[\tilde{n}]!} , \quad (1)$$

where  $q$  is positive,  $z$  is complex, and

$$\begin{aligned} [n]! &= [1][2] \dots [n] , & [k] &= 1 + q + q^2 + \dots + q^{k-1} , \\ [\tilde{n}]! &= [\tilde{1}][\tilde{2}] \dots [\tilde{n}] , & [\tilde{k}] &= 1 + \frac{1}{q} + \frac{1}{q^2} + \dots + \frac{1}{q^{k-1}} . \end{aligned} \quad (2)$$

Hence we immediately get  $E_q^z = e_{1/q}^z$ . Another, more popular, form of  $E_q^z$  is obtained using the identity

$$[\tilde{n}]! = q^{\frac{(1-n)n}{2}} [n]! . \quad (3)$$

Both exponential functions can be represented by infinite products,

$$e_q^z = \prod_{k=0}^{\infty} (1 - (1-q)q^k z)^{-1} , \quad E_q^z = \prod_{k=0}^{\infty} (1 + (1-q)q^k z) . \quad (4)$$

From this form we easily see that  $e_q^z E_q^{-z} = 1$ . Moreover,

$$D_q e_q^z = e_q^z, \quad D_q E_q^z = E_q^{qz}, \quad (5)$$

where  $D_q$  ( $q$ -derivative or Jackson's derivative) is defined by

$$D_q f(z) := \frac{f(qz) - f(z)}{qz - z}. \quad (6)$$

The existence of two representations of  $q$ -exponential functions (infinite series and infinite product) is related to well known formulae for the usual exponential function ( $q = 1$ ),

$$e^z = \sum_{k=0}^{\infty} \frac{z^k}{k!} = \lim_{m \rightarrow \infty} \left(1 + \frac{z}{m}\right)^m. \quad (7)$$

Two exponential functions of the quantum calculus generate two pairs of the  $q$ -trigonometric functions. Using notation of [20] we have:

$$\begin{aligned} \sin_q x &= \frac{e_q^{ix} - e_q^{-ix}}{2i}, & \text{Sin}_q x &= \frac{E_q^{ix} - E_q^{-ix}}{2i}, \\ \cos_q x &= \frac{e_q^{ix} + e_q^{-ix}}{2}, & \text{Cos}_q x &= \frac{E_q^{ix} + E_q^{-ix}}{2}. \end{aligned} \quad (8)$$

Taking into account properties of  $q$ -exponential functions (see above) we easily derive properties of standard  $q$ -trigonometric functions:

$$\cos_q x \text{Cos}_q x + \sin_q x \text{Sin}_q x = 1, \quad (9)$$

$$\sin_q x \text{Cos}_q x = \cos_q x \text{Sin}_q x,$$

$$\begin{aligned} D_q \sin_q x &= \cos_q x, & D_q \cos_q x &= -\sin_q x, \\ D_q \text{Sin}_q x &= \text{Cos}_q(qx), & D_q \text{Cos}_q x &= -\text{Sin}_q(qx). \end{aligned} \quad (10)$$

Note that the corresponding tangents coincide:  $\text{Tan}_q x = \tan_q x$ .

## 2 Improved $q$ -exponential function

New  $q$ -exponential function  $\mathcal{E}_q^z$  is defined as

$$\mathcal{E}_q^z := e_q^{\frac{z}{2}} E_q^{\frac{z}{2}} = \prod_{k=0}^{\infty} \frac{1 + q^k(1-q)\frac{z}{2}}{1 - q^k(1-q)\frac{z}{2}}, \quad (11)$$

where  $e_q^z, E_q^z$  are standard  $q$ -exponential functions. This definition is motivated by the classical Cayley transformation

$$z \rightarrow \text{cay}(z, a) := \frac{1 + az}{1 - az}, \quad (12)$$

see, e.g., [9, 17]. Indeed,

$$\mathcal{E}_q^{qz} = \frac{1 - (1-q)\frac{z}{2}}{1 + (1-q)\frac{z}{2}} \mathcal{E}_q^z = \text{cay}\left(-\frac{z}{2}, 1-q\right) \mathcal{E}_q^z. \quad (13)$$

**Theorem 1.** *The  $q$ -exponential function  $\mathcal{E}_q^z$  is analytic in the disc  $|z| < R_q$  and*

$$\mathcal{E}_q^z = \sum_{n=0}^{\infty} \frac{z^n}{\{n\}!}, \quad (14)$$

for  $|z| < R_q$ , where

$$R_q = \begin{cases} \frac{2}{1-q} & \text{for } 0 < q < 1, \\ \frac{2q}{q-1} & \text{for } q > 1, \\ \infty & \text{for } q = 1, \end{cases} \quad (15)$$

$$\{n\} := \frac{1 + q + \dots + q^{n-1}}{\frac{1}{2}(1 + q^{n-1})} = \frac{[n]}{\frac{1}{2}(1 + q^{n-1})} = \frac{2(1 - q^n)}{(1 - q)(1 + q^{n-1})}, \quad (16)$$

and, finally,  $\{n\}! = \{1\}\{2\} \dots \{n\}$ .

*Proof:* In the disc  $|z| < 1$  both series (1) are absolutely convergent for any  $q \in \mathbb{R}_+$ . Multiplying them we get

$$e_q^{\frac{z}{2}} E_q^{\frac{z}{2}} = \sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \frac{q^{\frac{j(j-1)}{2}} \left(\frac{z}{2}\right)^{k+j}}{[k]![j]!} = \sum_{n=0}^{\infty} \frac{\left(\frac{z}{2}\right)^n}{[n]!} \left( \sum_{j=0}^n \frac{q^{\frac{j(j-1)}{2}} [n]!}{[j]![n-j]!} \right). \quad (17)$$

Using Gauss's binomial formula (see, e.g., [20], formula (5.5))

$$\prod_{j=0}^{n-1} (z + a q^j) = \sum_{j=0}^n \frac{q^{\frac{j(j-1)}{2}} [n]!}{[j]![n-j]!} a^j z^{n-j}, \quad (18)$$

we have, as a particular case,

$$\sum_{j=0}^n \frac{q^{\frac{j(j-1)}{2}} [n]!}{[j]![n-j]!} = (1+1)(1+q) \dots (1+q^{n-1}). \quad (19)$$

Substituting (19) into (17) we get the formula (14) with  $\{n\}$  defined by (16). In order to obtain the radius of convergence, we compute

$$\lim_{n \rightarrow \infty} \left| \frac{z^{n+1}}{\{n+1\}!} \right| \left| \frac{\{n\}!}{z^n} \right| = \lim_{n \rightarrow \infty} \left| \frac{z}{\{n+1\}} \right| = \begin{cases} \frac{(1-q)|z|}{2} & \text{for } q < 1 \\ \frac{(q-1)|z|}{2q} & \text{for } q > 1 \end{cases} \quad (20)$$

Then, using d'Alembert's test, we get (for  $q \neq 1$ ) the radius of convergence (15). Note that  $R_{1/q} = R_q$ . In the case  $q = 1$  all  $q$ -exponential functions coincide with  $e^z$ , hence  $R_1 = \infty$ .  $\square$

**Theorem 2.** *The  $q$ -exponential function  $\mathcal{E}_q^z$  has the following properties:*

$$\mathcal{E}_q^{-z} = (\mathcal{E}_q^z)^{-1}, \quad |\mathcal{E}_q^{ix}| = 1, \quad (21)$$

$$\mathcal{E}_q^z = \mathcal{E}_{1/q}^z, \quad D_q \mathcal{E}_q^z = \langle \mathcal{E}_q^z \rangle, \quad (22)$$

where  $z \in \mathbb{C}$ ,  $x \in \mathbb{R}$  and we use the notation  $\langle f(z) \rangle := \frac{f(z) + f(qz)}{2}$ .

*Proof:* The first equation of (21) is a straightforward consequence of the definition (11). Then,  $\overline{\mathcal{E}_q^z} = \mathcal{E}_q^{\bar{z}}$ . Hence,

$$|\mathcal{E}_q^{ix}|^2 = \overline{\mathcal{E}_q^{ix}} \mathcal{E}_q^{ix} = \mathcal{E}_q^{-ix} \mathcal{E}_q^{ix} = 1. \quad (23)$$

The symbol  $\{n\}$  depends on  $q$ . In this proof it is convenient to use more precise notation  $\{n\} \equiv \{n\}_q$ ,  $\{n\}! \equiv \{n\}_q!$ . The equation  $\mathcal{E}_q^z = \mathcal{E}_{1/q}^z$  follows immediately from the obvious identity

$$\{n\}_q! = \{n\}_{1/q}! . \quad (24)$$

Finally,

$$D_q \mathcal{E}_q^z = \frac{\mathcal{E}_q^{qz} - \mathcal{E}_q^z}{qz - z} = \frac{\mathcal{E}_q^z}{(q-1)z} \left( \frac{1 - (1-q)\frac{z}{2}}{1 + (1-q)\frac{z}{2}} - 1 \right) = \frac{\mathcal{E}_q^{qz}}{1 + (1-q)\frac{z}{2}} , \quad (25)$$

$$\langle \mathcal{E}_q^z \rangle = \frac{1}{2} (\mathcal{E}_q^{qz} + \mathcal{E}_q^z) = \frac{1}{2} \left( \frac{1 - (1-q)\frac{z}{2}}{1 + (1-q)\frac{z}{2}} + 1 \right) \mathcal{E}_q^z = \frac{\mathcal{E}_q^{qz}}{1 + (1-q)\frac{z}{2}} , \quad (26)$$

which implies the second equation of (22).  $\square$

The properties (21) are identical with analogical properties of the exponential function  $e^z$ . We point out that neither  $e_q^z$  nor  $E_q^z$  satisfies (21). Instead, we have  $E_q^{-z} e_q^z = 1$ .

### 3 Improved $q$ -trigonometric functions

New  $q$ -sine and  $q$ -cosine functions are defined in a natural way:

$$\mathcal{S}in_q x = \frac{\mathcal{E}_q^{ix} - \mathcal{E}_q^{-ix}}{2i} , \quad \mathcal{C}os_q x = \frac{\mathcal{E}_q^{ix} + \mathcal{E}_q^{-ix}}{2} . \quad (27)$$

**Theorem 3.**  *$q$ -Trigonometric functions defined by (27) satisfy:*

$$\begin{aligned} \mathcal{C}os_q^2 x + \mathcal{S}in_q^2 x &= 1 , \\ D_q \mathcal{S}in_q x &= \langle \mathcal{C}os_q x \rangle , \\ D_q \mathcal{C}os_q x &= -\langle \mathcal{S}in_q x \rangle , \end{aligned} \quad (28)$$

*Proof:* Properties (28) follow directly from (21), (22) (note that  $\mathcal{E}_q^{ix} \mathcal{E}_q^{-ix} = 1$ ).  $\square$

**Corollary 4.**  *$q$ -Trigonometric functions  $\mathcal{C}os_q x$ ,  $\mathcal{S}in_q x$  are real for  $x \in \mathbb{R}$ . Moreover, for any  $x \in \mathbb{R}$ , we have*

$$-1 \leq \mathcal{C}os_q x \leq 1 , \quad -1 \leq \mathcal{S}in_q x \leq 1 . \quad (29)$$

**Theorem 5.** *New  $q$ -trigonometric functions can be expressed by standard  $q$ -trigonometric functions as follows:*

$$\begin{aligned}\mathcal{C}os_q 2x &= \cos_q x \mathcal{C}os_q x - \sin_q x \mathcal{S}in_q x = \frac{1 - \tan_q^2 x}{1 + \tan_q^2 x} , \\ \mathcal{S}in_q 2x &= \sin_q x \mathcal{C}os_q x + \cos_q x \mathcal{S}in_q x = \frac{2 \tan_q x}{1 + \tan_q^2 x} .\end{aligned}\tag{30}$$

*Proof:* First, we compute

$$\begin{aligned}\cos_q x \mathcal{C}os_q x - \sin_q x \mathcal{S}in_q x &= \frac{e_q^{ix} E_q^{ix} + e_q^{-ix} E_q^{-ix}}{2} = \mathcal{C}os_q 2x , \\ \sin_q x \mathcal{C}os_q x + \cos_q x \mathcal{S}in_q x &= \frac{e_q^{ix} E_q^{ix} - e_q^{-ix} E_q^{-ix}}{2} = \mathcal{S}in_q 2x .\end{aligned}\tag{31}$$

Then, using (9), we get

$$\begin{aligned}\mathcal{C}os_q 2x &= \frac{\cos_q x \mathcal{C}os_q x - \sin_q x \mathcal{S}in_q x}{\cos_q x \mathcal{C}os_q x + \sin_q x \mathcal{S}in_q x} = \frac{1 - \tan_q x \mathcal{T}an_q x}{1 + \tan_q x \mathcal{T}an_q x} , \\ \mathcal{S}in_q 2x &= \frac{\sin_q x \mathcal{C}os_q x + \cos_q x \mathcal{S}in_q x}{\cos_q x \mathcal{C}os_q x + \sin_q x \mathcal{S}in_q x} = \frac{\tan_q x + \mathcal{T}an_q x}{1 + \tan_q x \mathcal{T}an_q x} .\end{aligned}\tag{32}$$

Taking into account  $\mathcal{T}an_q x = \tan_q x$  we complete the proof.  $\square$

## 4 Conclusions

Motivated by the classical Cayley transformation and recent results in the time scales calculus (see [9]), we introduced a new definition of the  $q$ -exponential function. Main advantages of the new  $q$ -exponential function consist in better qualitative properties (i.e., its properties are more similar to properties of  $e^z$ ). In particular, it maps the unitary axis into the unit circle, compare (21), which implies excellent properties of new trigonometric functions, including formulae (28) and boundedness (29).

Especially interesting is the Pythagorean identity:  $\mathcal{C}os_q^2 x + \mathcal{S}in_q^2 x = 1$ . According to our best knowledge, other  $q$ -defomations of trigonometric functions do not satisfy this property. The same concerns even the paper [15], full of surprising identities.

Our exponential function is closely related to both popular  $q$ -exponential functions (1). Therefore, proofs and calculations concerning  $\mathcal{E}_q^z$  can be usually done with the help of known results. We plan to express in terms of the new exponential function classical results containing  $q$ -exponential functions, and we hope to obtain some improvements.

## References

- [1] G.E.Andrews, R.Askey, R.Roy: *Special functions*, Cambridge Univ. Press 1999.
- [2] N.M.Atakishiyev: “On a one-parameter family of  $q$ -exponential functions”, *J. Phys. A: Math. Gen.* **29** (1996) L223-L227.
- [3] N.M.Atakishiyev: “On the Fourier-Gauss transforms of some  $q$ -exponential and  $q$ -trigonometric functions”, *J. Phys. A: Math. Gen.* **29** (1996) 7177-7181.
- [4] N.M.Atakishiyev, A.U.Klimyk, K.B.Wolf: “A discrete quantum model of the harmonic oscillator”, *J. Phys. A: Math. Theor.* **41** (2008) 085201 (14pp).
- [5] L.C.Biedenharn: “The quantum group  $SU_q(2)$  and a  $q$ -analogue of the boson operators”, *J. Phys. A: Math. Gen.* **22** (1989) L873-L878.
- [6] M.Bohner, G.S.Guseinov: “The  $q$ -Laplace and  $h$ -Laplace transforms”, *J. Math. Anal. Appl.* **365** (2010) 75-92.
- [7] J.Bustoz, J.L.Cardoso: “Basic analog of Fourier series on a  $q$ -linear grid”, *J. Approx. Theory* **112** (2001) 134-157.
- [8] I.Cherednik: “Whittaker limits of difference spherical functions”, *Int. Math. Res. Notices* **2009**(2009) 3793-3842.
- [9] J.L.Cieřliński: “New definitions of exponential, hyperbolic and trigonometric functions on time scales”, *preprint arXiv: 1003.0697 [math.CA]* (2009).
- [10] R.Díaz, E.Pariguán: “An example of Feynman-Jackson integrals”, *J. Phys. A: Math. Theor.* **40** (2007) 1265-1272.
- [11] T.Ernst: “The history of  $q$ -calculus and a new method”, UUDM Report 2000:16, Dept. Math. Uppsala Univ. 2000.
- [12] K.Ey, A.Ruffing, S.Suslov: “Method of separation of the variables for basic analogs of equations of mathematical physics”, *Ramanujan J.* **13** (2007) 407-447.
- [13] R.J.Finkelstein: “ $q$ -Uncertainty relations”, *Int. J. Mod. Phys. A* **13** (1998) 1795-1803.
- [14] G.Gasper, M.Rahman: *Basic hypergeometric series*, Cambridge Univ. Press 1990.
- [15] R.Wm.Gasper: “Experiments and discoveries in  $q$ -trigonometry”, [in:] *Symbolic computation, number theory, special functions, physics and combinatorics*, pp. 79-105; edited by F.G.Garvan and M.E.H.Ismail, Kluwer Academic Publishers 2001.
- [16] S.Hilger: “Analysis on measure chains – a unified approach to continuous and discrete calculus”, *Results Math.* **18** (1990) 19-56.
- [17] A.Iserles: “On Cayley-transform methods for the discretization of Lie-group equations”, *Found. Comput. Math.* **1** (2001) 129-160.
- [18] F.H.Jackson: “A basic-sine and cosine with symbolical solutions of certain differential equations”, *Proc. Edin. Math. Soc.* **22** (1904) 28-38.



- [19] F.H.Jackson: "On  $q$ -difference equations", *Am. J. Math.* **32** (1910) 305-314.
- [20] V.Kac, P.Cheung: *Quantum calculus*, Springer 2002.
- [21] B.A.Kupershmidt: " $q$ -Newton binomial: from Euler to Gauss", *J. Nonlin. Math. Phys.* **7** (2000) 244-262.
- [22] A.K.Kwaśniewski, B.K.Kwaśniewski: "On  $q$ -difference equations and  $Z_n$  decompositions of  $\exp_q$  function", *Adv. Appl. Clifford Alg.* **11** (2001) 39-61.
- [23] A.Lorek, A.Ruffing, J.Weiss: "A  $q$ -deformation of the harmonic oscillator", *Z. Phys. C* **74** (1997) 369-377.
- [24] D.S.McAnally: " $q$ -Exponential and  $q$ -gamma functions. I.  $q$ -Exponential functions", *J. Math. Phys.* **36** 546-573.
- [25] A.J.Macfarlane: "On  $q$ -analogues of the quantum harmonic oscillator and the quantum group  $SU(2)_q$ ", *J. Phys. A: Math. Gen.* **22** (1989) 4581-4588.
- [26] C.Quesne: "Disentangling  $q$ -exponentials: a general approach", *Int. J. Theor. Phys.* **43** 545-559.
- [27] S.K.Suslov: " "Addition" theorems for some  $q$ -exponential and  $q$ -trigonometric functions", *Meth. Appl. Anal.* **4** (1997) 11-32.
- [28] S.K.Suslov: "Another addition theorem for the  $q$ -exponential function", *J. Phys. A: Math. Gen.* **33** (2000) L375-L380.
- [29] M.R.Ubriaco: "Time evolution in quantum mechanics on the quantum line", *Phys. Lett. A* **163** (1992) 1-4.